





Percentage of Penicillin Non-Susceptible *S. pneumoniae* in Canada: 1988-2009



*Non-meningitis, Oral pen breakpoints used




Percentage of Penicillin* Non-Susceptible *S. pneumoniae* in Canada: 1993-2009



*Non-meningitis, IV breakpoints used

Canadian Bacterial Surveillance Network, Jun 2010


Percentage of Non-susceptible Sterile Site vs Non-sterile Site Isolates of *S. pneumoniae*, 2009



*Non-meningitis breakpoints used, Non-meningitis (IV) for pen




Percentage of Amoxicillin Non-Susceptible *S. pneumoniae* in Canada: 1988-2009






Percentage of Non-susceptible Isolates of *S. pneumoniae* in Ages ≤ 15 and >64 , 2009



*Non-meningitis breakpoints used, Non-meningitis (IV) for pen




Percentage of Non-susceptible Isolates of *S. pneumoniae* in Geographic Regions of Canada, 2009



*Non-meningitis breakpoints used




Macrolide-Resistant Pneumococci: Canadian Bacterial Surveillance Network, 1993-2009







Fluoroquinolone-Resistant Pneumococci: Canadian Bacterial Surveillance Network, 1997-2009






Fluoroquinolone-Resistant Pneumococci in Respiratory Isolates from Adults >64 years: 1988-2009





Serotype distribution by year: 1994-2009



■ PCV7 ■ PCV10 additional serotypes (1,5,7F) ■ PCV13 additional serotypes (3,6A,19A) ■ NPCV



Serotype distribution by year: 1994-2009

